

Evaluation of the Relationship Between Level of Nicotine Dependence and Risk of Anxiety, Depression and Some Socio-Demographic Variables

Yasemin SAĞLAN^{1,a}, Hüseyin BALCIOĞLU¹

¹Eskişehir Osmangazi University Faculty of Medicine, Department of Family Medicine, Eskişehir, Türkiye

ABSTRACT

Objective: Nicotine addiction remains the leading cause of preventable disease and premature death in many countries. The aim of this study is to determine the frequency of high levels of nicotine dependence and related factors in patients admitted to the smoking cessation clinic.

Material and Method: The study is a cross-sectional study conducted on patients who applied to Smoking Cessation Polyclinic affiliated with a district health department. The Hospital Anxiety Depression Scale, which is a screening scale, was used in the psychological evaluation of individuals, and the Fagerström Nicotine Dependence Test was used to evaluate nicotine addiction. Logistic regression analysis was performed to determine the independent variables associated with nicotine addiction.

Results: The study group consisted of a total of 318 patients. 60.4 % of the study group was male. The age of the study group ranged from 19 to 70 years. In the study, the frequency of those with high levels of nicotine addiction was found to be 25.8 % (n =82) . In a multivariate analysis, the risk of high levels of nicotine addiction was higher in men, singles, smokers with tea or coffee and those with anxiety (p <0.05).

Conclusion: Nicotine dependence is an important public health problem. In patients with anxiety, it was detected 10.154 times higher than those without high level of nicotine dependence.

Keywords: Nicotine, Anxiety, Depression, Addiction.

ÖZ

Nikotin Bağımlılığı Düzeyi ile Anksiyete, Depresyon Riski ve Bazı Sosyo-Demografik Değişkenler Arasındaki İlişkinin Değerlendirilmesi

Amaç: Nikotin bağımlılığı birçok ülkede önlenebilir hastalıkların ve erken ölümün önde gelen nedeni olmaya devam etmektedir. Bu çalışmanın amacı sigara bırakma polikliniğine başvuran hastalarda yüksek düzeyde nikotin bağımlılığı sıklığını ve ilişkili faktörleri belirlemektir.

Gereç ve Yöntem: Araştırma bir ilçe sağlık müdürlüğüne bağlı Sigara Bırakma Polikliniğine başvuran hastalar üzerinde gerçekleştirilen kesitsel bir çalışmındır. Bireylerin psikolojik değerlendirilmesinde tarama ölçüği olan Hastane Anksiyete Depresyon Ölçeği, nikotin bağımlılığının değerlendirilmesinde ise Fagerström Nikotin Bağımlılığı Testi kullanıldı. Nikotin bağımlılığı ile ilişkili bağımsız değişkenleri belirlemek için lojistik regresyon analizi yapıldı.

Bulgular: Çalışma grubu toplam 318 hastadan oluşmaktadır. Çalışma grubunun %60,4' ü erkektir. Çalışma grubunun yaşları 19 ile 70 arasında değişmektedir. Araştırmada nikotin bağımlılığı yüksek olanların sıklığı %25,8 (n =82) olarak belirlendi. Çok değişkenli bir analizde yüksek düzeyde nikotin bağımlılığı riski erkeklerde, bekârlarda, çay veya kahve içenlerde ve anksiyetesine olanlarda daha yüksekti (p <0,05).

Sonuç: Nikotin bağımlılığı önemli bir halk sağlığı sorunudur. Anksiyetiesi olan hastalarda nikotin bağımlılığı yüksek olmayanlara göre 10.154 kat daha fazla tespit edildi.

Anahtar Sözcükler: Nikotin, Anksiyete, Depresyon, Bağımlılık.

Bu makale atıfta nasıl kullanılır: Sağlan Y, Balcioglu H. Nikotin Bağımlılığı Düzeyi ile Anksiyete, Depresyon Riski ve Bazı Sosyo-Demografik Değişkenler Arasındaki İlişkinin Değerlendirilmesi. Fırat Tip Dergisi 2025; 30(4): 268-273.

How to cite this article: Saglan Y, Balcioglu H. Evaluation of the Relationship Between Level of Nicotine Dependence and Risk of Anxiety, Depression and Some Socio-Demographic Variables. Fırat Med J 2025; 30(4): 268-273.

ORCID IDs: Y.S. 0000-0002-5066-0432, H.B. 0000-0003-1648-3206.

Tobacco use is a common public health problem in our country as well as all over the world. It causes fatal health problems due to direct or passive smoke exposure. Tobacco use is decreasing in developed countries and increasing in underdeveloped and developing countries. It is predicted that 80 % of tobacco-related deaths will occur in developing societies by 2030 (1, 2).

Smokers are not aware that it is an addiction. He thinks it is a habit, but cigarettes are an addictive substance with nicotine. Although a significant portion of smo-

kers are not satisfied with smoking, they continue their smoking behavior due to nicotine addiction. Important studies are carried out in our country in the fight against tobacco, which has a higher prevalence of addiction than any other substance at any age. One of the studies in the global fight against addiction is smoking cessation polyclinics. In our country, the ban on smoking indoors, the social awareness created through the media, the diseases it causes and economic reasons have increased the demands for smoking cessation outpatient clinics (3-5).

*Yazışma Adresi: Yasemin SAĞLAN, Eskişehir Osmangazi University Faculty of Medicine, Department of Family Medicine, Eskişehir, Türkiye
Tel: 0222 239 29 79

Geliş Tarihi/Received: 13.06.2024

* This study was presented as a oral presentation at 4th International Congress on Medical and Health Sciences Research (22-23 rd August2020, Hattuşaş).

Studies have shown that the probability of quitting without the support of a smoking cessation outpatient clinic is 3-5 %, and the rate of quitting with outpatient support is 40 % (6). Smoking cessation treatment should be comprehensive and continuous, as should the approach to substance abuse. This comprehensive evaluation should include many factors such as the person's sociodemographic information, medical history, smoking frequency and psychological evaluation. Examination of these factors in interviews to be held in smoking cessation outpatient clinics will guide the determination of the treatment method.

It is known that approximately half of the patients with psychiatric disorders in our country and 40-90 % of the psychiatric patients are cigarette addicts (7). The American Psychiatric Association defines tobacco habit as a psychiatric disorder that includes cognitive, behavioral and physiological symptoms (8). Therefore, the treatment of a smoker should be similar to the approach to substance abuse. The effectiveness of the service offered to voluntary individuals who want to quit smoking increases in proportion to the extensive interviews. The Nicotine Addiction Level and Anxiety Depression Level to be determined are important factors in shaping the treatment.

The treatment of a smoker should be like the approach to substance abuse. The effectiveness of the service offered to voluntary individuals who want to quit smoking increases in proportion to the comprehensive interviews. The aim of this study is to determine the frequency of high levels of nicotine dependence and related factors in patients admitted to the smoking cessation clinic. In addition, it is to evaluate the relationship between the level of nicotine addiction and the risk of anxiety and depression.

MATERIAL AND METHOD

The study is a cross-sectional study conducted on patients who applied to Smoking Cessation Polyclinic affiliated with a district health department between September 2020 and January 2021. When the smoking frequency was accepted as 27 %, the margin of error was 5 % and the confidence interval was 95 %, the minimum number of patients to be reached was calculated as 302. A total of 318 patients who agreed to participate constituted the study group. Volunteer patients who applied to Smoking Cessation Clinic for the first time were included in the study. In our study, the

sociodemographic characteristics and detailed smoking history of the participants were recorded in the forms by face-to-face interview method. Age grouping was based on age 45 because there is a rapid increase in smoking cessation rates at age 45 (9). Detailed smoking history includes questions such as 'The reasons that increase the desire to smoke' and 'The reason for starting smoking'. Frequently seen reasons from the literature were taken as basis when creating question options (10-14).

The individuals participating in the study read and approved the Informed Consent Form. The Fagerström Nicotine Dependence Test (FNBT), which is the most common method, was used to evaluate the nicotine addiction level of individuals. FNBT was developed by Fagerstrom et al. (15) and consists of 6 questions. The Turkish validity and reliability study of the Fagerstrom Nicotine Dependence Test was conducted by Uysal et al. in 2004 (16). The Turkish version of FTND had moderate reliability (Cronbach alpha: 0.56). Each question has certain points according to the answer. It is evaluated as low (0 - 4 points); medium (5 - 6 points); high (7 - 8 points) and very high (9 - 10 points).

The Hospital Anxiety Depression Scale (HAD), a screening scale, was used in the psychological evaluation of individuals. HAD was developed by Zigmond and Snaith in 1983 (17). The Turkish validity and reliability study performed by Aydemir et al. (18) and the Cronbach alpha coefficient was found to be 0.8525 for the anxiety subscale and 0.7784 for the depression subscale. HAD contains a total of 14 questions, 7 of which measure anxiety and 7 of which measure depression. In the Turkish version of the scale, the cut-off point for the anxiety subscale (HAD-A) was determined as 10, and the depression subscale (HAD-D) was determined as 7. Accordingly, those who score above these points are considered at risk.

Data were analyzed in the IBM SPSS (version 21.0) statistical package program. Continuous variables are given as mean \pm standard deviation, and categorical variables are given as frequency and percentage. Logistic regression analysis was performed to identify independent variables associated with nicotine addiction.

RESULTS

Sixty-four% of the study group was male. The age of the study group ranged from 19 to 70, with a mean (SD) of 40.10 (10.63) years (Table 1).

Table 1. The sociodemographic attributes and factors associated with nicotine dependence of the study group.

| Sociodemographic characteristics and factors associated with nicotine dependence | | Number (n) | Percentage (%) |
|--|--|------------|----------------|
| Gender | | | |
| Female | | 126 | 39.6 |
| Male | | 192 | 60.4 |
| Age group | | | |
| <45 | | 216 | 67.9 |
| ≥45 | | 102 | 32.1 |
| Education status | | | |
| Primary education | | 128 | 40.3 |
| High school education or higher | | 190 | 59.7 |
| Marital status | | | |
| Married | | 234 | 73.6 |
| Not married | | 84 | 26.4 |
| The reasons that increase the desire to smoke | | | |
| Stress | | 124 | 39.0 |
| Postprandial | | 88 | 27.7 |
| Tea/Coffee | | 106 | 33.3 |
| The reason for applying to the smoking cessation clinic | | | |
| Doctor's recommendation | | 46 | 14.5 |
| Voluntarily | | 272 | 85.5 |
| The presence of another smoker at home | | | |
| No | | 172 | 54.1 |
| Yes | | 146 | 45.9 |
| The presence of a smoker at work | | | |
| No | | 116 | 36.5 |
| Yes | | 202 | 63.5 |
| The reason for starting smoking | | | |
| Curiosity | | 92 | 28.9 |
| Wannabe | | 144 | 45.3 |
| Stress/Worry | | 40 | 12.6 |
| The others | | 42 | 13.2 |
| Hospital Anxiety | | | |
| No | | 193 | 60.7 |
| Yes | | 125 | 39.3 |
| Hospital Depression | | | |
| No | | 165 | 51.9 |
| Yes | | 153 | 48.1 |

When the factors related to nicotine addiction of the individuals in our study were examined, it was determined that 60.4 % were male, 67.9 % (n =216) were under 45 years old, 73.6 % were married (n =234), 39.0 % (n =124) of them were stress and 85.5 % (n =272) of the applications to the smoking cessation outpatient clinic were voluntary.

In the study group, anxiety was found in 39.3% (n=125) and depression in 48.1% (n=153).

The frequency of those with a high level of nicotine dependence was found to be 25.8 % (n =82).

In a multivariate analysis, the risk of high level of nicotine dependence were higher among male (OR: 2.360), unmarried (OR: 1.942), in those who smoked alongside tea or coffee (OR: 2.219), in those with anxiety (OR: 10.154) (for each; $p <0.05$).

Additionally, no relationship was found between nicotine addiction level and age, educational status, the reason for applying to the smoking cessation clinic, the presence of another smoker at home, the presence of a smoker at work, the reason for starting smoking, the type of treatment, depression (Table 2).

Table 2. Multivariate logistic regression analysis outcomes.

| Variables | High Levels of Nicotine Dependence(%) | OR (95% CI) | p |
|--|---------------------------------------|-------------|------------------|
| Gender | Female | 19.0 | 0.011 |
| | Male | 30.2 | |
| Marital status | Married | 21.4 | 0.042 |
| | Not married | 38.1 | |
| The reasons that increase the desire to smoke | Stress | 19.4 | 0.072 |
| | Postprandial | 20.5 | |
| | Tea/Coffee | 37.7 | |
| Hospital Anxiety | No | 10.4 | <0.001 |
| | Yes | 9.6 | |
| Hospital Depression | No | 15.8 | 0.804 |
| | Yes | 36.6 | |

OR: Odd's ratio, CI: Confidence interval.

DISCUSSION

Smoking addiction is a condition that requires long-term follow-up and struggle. It is known that psychiatric disorders such as lifelong mood disorders, anxiety disorders, substance abuse and personality disorders are more common in smokers (19). Psychological disorders accompanying cigarette addiction affect the success of treatment, and it would be beneficial to evaluate it as a risk factor. All psychiatric illnesses are serious risk factors for nicotine addiction. Smoking cessation outpatient clinics play an important role in the follow-up and treatment of this addiction. There is no current study evaluating the interviews in the Healthy Life Centers Smoking Cessation Polyclinics in our province.

In our study, it was observed that the rate of admission to the smoking cessation outpatient clinic was higher in males with a rate of 60.4 %. Similarly, in many studies, men are more likely to apply to smoking cessation outpatient clinics. We can explain this situation with the high rate of smoking among men in our country (20-22).

When we look at the reasons that increase the desire to smoke in our study, it was seen that the most 39.0 % (n =124) of them was stress. In a study conducted with Medical Faculty students in our country, stress factor was found to be the reason for starting smoking at a rate of 38.3 %, similar to our study (23). In another study conducted in our country, the causes that increase the desire to smoke the most were found to be post-meal and stress (24). The serotonergic system is responsible for the etiology of both anxiety and depression, as well as addiction. 5-HT1A receptors play a role in coping with depression, anxiety, and stress, and 5-HT1B receptors play a role in substance addiction. It has been reported that irregularities in this system cause mood disorders, substance addiction, and suicide (25). In addition, studies have reported that addicts generally use emotion-focused and avoidant coping methods in stress management. In these methods, the person resorts to substance use to stay away from stress and to escape from problems. Tolerance develops over time, causing addiction (26, 27).

It was determined that 85.5 % (n =272) of the applications to the smoking cessation outpatient clinic were voluntary. In the study conducted by Özsahin et al. (28), it was found that approximately 20 % of women and 10 % of men quit smoking voluntarily. In our study, we think that there were applications due to social media or illness, but this was at high rates because individuals voluntarily expressed this situation.

In a multivariate analysis, we found that the risk of high-level nicotine addiction is 2.360 times greater in men than in women. Similarly, in the study conducted by Esen et al. (29), the level of addiction was found to be high in 60.40 % of men. In another study conducted in India, a high level of addiction was observed in men (30).

In a multivariate analysis, we found that the risk of high-level nicotine addiction was 1.942 times higher in singles. Again, in a large population-based study conducted on Nicobar and Andamar Islands, nicotine addiction rates were found to be higher in divorced individuals (30). In a study conducted in our country by Şahbaz et al. (31), it was found that smoking cessation rates were higher in married individuals than in single individuals. It is known that partner support during the smoking cessation process increases quitting rates. The lack of partner support during the smoking cessation process in single individuals may be the reason why nicotine addiction rates are higher than in married individuals (32, 33).

We did not find a significant relationship between the level of nicotine addiction and age and educational status. The data of a similar study conducted in the smoking cessation outpatient clinic in our country supports our study (29). Orsel et al. (34) in another study, it was found that there was a significant difference between cigarette addiction and age groups.

When we evaluate to the reasons that increase the desire to smoke in a multivariate analysis, we found that the risk of high-level nicotine addiction was 2.219 times higher in those who drink tea/coffee than stress and postprandial. There are publications in the literature that support our study and that smoking rates are higher after drinking tea. (10, 11, 35). In the study conducted by Kanik et al. (36), it was determined that individuals with coffee drinking habits had a significantly higher nicotine addiction level, and there was no significance between tea drinking habits and nicotine addiction levels. Drinking tea/coffee, which is one of the activities that accompanies smoking, is identified with smoking. The presence of one of these substances taken together causes a desire for the other (37, 38).

In our study, we found that the risk of high level of nicotine addiction was 10.154 times higher in those with anxiety. There are studies that support our study and found that anxiety is higher in smokers compared to non-smokers. It has been reported that anxiety is more effective than depression in cigarette addicts (39). Dorner et al. (40) reported that depression and anxiety were associated with high smoking frequency, high level of addiction and increased nicotine withdrawal complaints. In support of this study, the risk of nicotine addiction was not found to be significantly high in patients with depression in our study. In a meta-analysis study, it was determined that the risk of depression increased approximately 2 times in smokers compared to non-smokers or people who quit smoking (41). In another study, in which the relationship between smoking addiction and depression and anxiety was tried to be determined, the depression level was determined as 22.44 ± 13.54 and the anxiety level as 19.7 ± 12.5 (42). In the study conducted by McClave et al. (43) with cigarette addicts, it was reported that 20.3 % experienced lifelong depression, 14.9 % experienced lifelong anxiety, and 10.1 % experienced both depression and anxiety. Caykara et al. (44) when compared

according to gender, it was shown that there was a very high significant difference between anxiety levels and a high significant difference between depression levels.

Conclusion

The developing social awareness and awareness in our country increases the applications of smokers to smoking cessation outpatient clinics. When we evaluate these applications, anxiety and depression appear as risk factors in individuals with cigarette addiction. For this reason, it will be more effective to carry out medi-

cal treatment, psychosocial support and rehabilitation services together in the treatment of smoking addiction. Preventive studies should be conducted to address the causes of anxiety and rehabilitation and social support services should be provided to reduce anxiety levels which has been identified as a risk factor for high levels of smoking addiction.

Acknowledgement

No financial support was received.

REFERENCES

1. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006; 3: 442.
2. The W. Curbing the epidemic: governments and the economics of tobacco control. The World Bank. *Tob Control* 1999; 8: 196-201.
3. Kanıt L, Keser A. Tütün bağımlılığının biyofizyolojisi. Aytemur ZA, Akcay Ş, Elbek O, (Editörler). *Tütün ve Tütün Kontrolü*. Ankara: Türk Toraks Derneği Yayımları 2010; 10: 141-56.
4. Benowitz LN, Hukkanen J, Peyton J. Nicotine chemistry, metabolism, kinetics and biomarkers. Henningfield JE, London ED, Pogun S (Editors). *Nicotine psychopharmacology*. 192th edition 2009; 29-60.
5. Argüder E, Karalezli A, Hezer H ve ark. Sigara bırakma başarısını etkileyen faktörler. *Türk Toraks Dergisi* 2013; 14: 81-7.
6. Karlikaya C, Öztuna F, Aytemur Solak Z ve ark. Tütün kontrolü. *Toraks Dergisi* 2006; 7: 51-64.
7. Ertekin H, Aydin M, Arslan M et al. Are psychiatric outpatients assessed for smoking?. *Konuralp Med J* 2017; 9: 155-9.
8. Hughes JR, Fiester S, Goldstein MG et al. Practice guideline for the treatment of patients with nicotine dependence. *Am J Psychiatry* 1996; 153: 1-31.
9. DSÖ Küresel Yetişkin Tütün Araştırma Raporu www.who.int/tobacco/surveillance/survey/gats/report_tur_2012.
10. İtil O, Ergör G, Ceylan E. Knowledge and attitudes about smoking among students in a medical faculty. *Turkish Respirat J* 2004; 5: 86-91.
11. Ünüvar EM, Dişçigil G. Sigarayı bırakma girişimlerinde başarayı etkileyen faktörler-hekim adayları örneği. *J Turkish Family Physic* 2017; 8: 57-65.
12. Aydoğan ES. İstanbul'da genel bir devlet hastanesinde çalışanların sigara içme durumu ve etkileyen faktörler. *İstanbul Med J* 2013; 14: 170-4.
13. Atilgan Y, Gürkan S, Şen E. Hastanemizde çalışan personelin sigara içme durumu ve etkileyen faktörler. *Türk Toraks Dergisi* 2008; 9: 160-6.
14. Yakar B, Pirinçci E, Kurt Canpolat E. Sigara İçme Durumuna Etki Eden Faktörler ve Tütün Kontrol Eylem Planının Sigara Kullanım Üzerine Etkisinin Araştırılması. *Fırat Üniversitesi Sağlık Bilimleri Tıp Dergisi* 2020; 34: 147-153.
15. Heatherton TF, Kozlowski LT, Frecker RC et al. The Fagerström test for nicotine dependence: a revision of the Fagerstrom Tolerance Questionnaire. *British J Add* 1991; 86: 1119-27.
16. Uysal MA, Kadakal F, Karşıdağ C et al. Fagerström test for nicotine dependence: reliability in a Turkish sample and factor analysis. *Tuberk Toraks* 2004; 52: 115-21.
17. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica* 1983; 67: 361-70.
18. Aydemir OJTPD. Hastane anksiyete ve depresyon ölçeği Türkçe formunun geçerlilik ve güvenilirliği. *Türk Psikiyatri Derg* 1997; 8: 187-280.
19. Black DW, Zimmerman M, Coryell WH. Cigarette smoking and psychiatric disorder in a community sample. *Ann Clin Psych* 1999; 11: 129-36.
20. Godtfredsen NS, Lam TH, Hansel TT et al. COPD-related morbidity and mortality after smoking cessation: status of the evidence. *Eur Resp J* 2008; 32: 844-53.
21. Barbano G, Bressan MC, Nardini S. The stage-of-change model in smoking cessation in respiratory patients: does it need to be revisited. *Eur Resp Soc J* 2008; 42: 51-6.

22. Küresel Yetişkin Tütün Araştırması. Türkiye İstatistik Kurumu 2012. http://www.tuik.gov.tr/PreHaber_Bultenleri.
23. Er T, Kurçer MA. Tıp Fakültesi 1. ve 6. Sınıf Öğrencilerinin Sigara İçme Davranışları ve Anksiyete Düzeyleri. Bağımlılık Dergisi 2020; 21: 201-9.
24. Uzaslan EK, Ozyardimci N, Karadag MA et al. The physician's intervention in smoking cessation: Results of the five years of smoking cessation clinic. Ann Med Sci 2000; 9: 63-9.
25. Tamam L, Zeren T. Depresyonda serotonerjik düzenekler. Klinik Psikiyatri Dergisi 2002; 5: 11-8.
26. Cooper ML, Russell M, Skinner JB et al. Stress and alcohol use: moderating effects of gender, coping, and alcohol expectancies. J Abnormal Psychol 1992; 101: 139.
27. Simons RL, Robertson JF. The impact of parenting factors, deviant peers, and coping style upon adolescent drug use. Family Relations 1989; 27: 38-1.
28. Ozsahin K, Unsal A, Erdogan F ve ark. Sigara bırakma üzerinde etkili faktörler: Aile hekimliği poliklinik hastaları üzerinde bir çalışma. TAF Prev Med Bull 2007; 6: 181-6.
29. Esen AD, Arica S. Sigarayı bırakmak için başvurularda nikotin bağımlılık düzeyi ve sosyodemografik özelliklerin değerlendirilmesi. Ankara Med J 2018; 18: 328-36.
30. Manimunda SP, Benegal V, Sugunan AP et al. Tobacco use and nicotine dependency in a cross-sectional representative sample of 18,018 individuals in Andaman and Nicobar Islands, India. BMC Public Health 2012; 12: 1-11.
31. Ahbaz S, Kılınç O, Günay T ve ark. Sigara İçme ve Demografik Özelliklerin Sigara Bırakma Tedavilerinin Sonuçlarına Etkileri. Toraks Dergisi 2007; 8: 110-4.
32. West R, McEwen A, Bolling K et al. Smoking cessation and smoking patterns in the general population: a 1-year follow-up. Addiction 2001; 96: 891-902.
33. Chandola T, Head J, Bartley M. Socio-demographic predictors of quitting smoking: how important are household factors?. Addiction 2004; 99: 770-7.
34. Örsel O, Örsel S, Alpar S et al. Sigara bırakmada nikotin bağımlılık düzeylerinin tedavi sonuçlarına etkisi. Solunum Hastalıkları 2005; 16: 112-8.
35. Fırat M, Ayran G, Kanbay Y. Sigara Bırakma Polikliniğine Başvuran Bireylerin Sigaraya Başlama, Sigara Kullanma ve Sigara Bırakmayla İlgili Özellikleri. Addicta: Turkish J Addic 2019; 6: 244-56
36. Kanık MK, Tözün M. Smoking and Other Tobacco Use Characteristics and Nicotine Dependence Status of Those Who Applied to Smoking Cessation Polyclinics in Izmir. Estüdam Halk Sağlığı Dergisi 2020; 5: 84-95.
37. Uzaslan E. Sigarayı bırakma yöntemleri. Sted J Cont Med Educ 2003; 12: 166-71.
38. Fidancı İ, Arslan İ, Tekin O ve ark. Sigara Bırakma Başarısında; Çay, Kahve, Alkol İçme Alışkanlıklarının, Kilo Alma Korkusunun ve Tedavi Yöntemlerinin Rolü. Konuralp Tıp Dergisi 2016; 8: 2.
39. Mykletun A, Overland S, Aarø LE et al. Smoking in relation to anxiety and depression: evidence from a large population survey: The HUNT study. Eur Psych 2008; 23: 77-84.
40. Dorner TE, Tröstl A, Womastek I et al. Predictors of short-term success in smoking cessation in relation to attendance at a smoking cessation program. Nicotine & Tobacco Res 2011; 13: 1068-75.
41. Luger TM, Suls J, Vander Weg MW. How robust is the association between smoking and depression in adults? A meta-analysis using linear mixed-effects models. Addictive Behaviors 2014; 39: 1418-29.
42. McCabe RE, Chudzik SM, Antony MM et al. Smoking behaviors across anxiety disorders. J Anx Dis 2004; 18: 7-18.
43. McClave AK, Dube SR, Strine TW et al. Associations between smoking cessation and anxiety and depression among US adults. Addictive Behaviors 2009; 34: 491-7.
44. Çaykara B, Tuna R, Sağlam ZA ve ark. Sigara bağımlılığında depresyon, umutsuzluk ve anksiyete düzeyleri arasındaki ilişkinin belirlenmesi. İzmir Göğüs Hastanesi Dergisi 2019; 33: 15-23.